

430 Rec'd PCT/PTO 28 JUL 1999

09/355296

1103326-0576

TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED OFFICE (DO/US)

PCT/SE99/00979 07 June 1999 09 June 1998 and 17 July 1998
International Application Number International Filing Date Priority Date(s) Claimed

RNA POLYMERASE ASSAY

Title of Invention

BALGANESH, Tanjore; RAMACHANDRAN, Vasanthi; and SHARMA, Umender

Applicant(s) for DO/US

| |
|---|
| <p>"Express Mail" Label No <u>EE040967375US</u></p> <p>Date of Deposit <u>JULY 28</u>, 1999.</p> <p>I hereby certify that this paper is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the: Assistant Commissioner for Patents, Washington, D.C. 20231.</p> <p><u>Mirella Ponce</u> (Type or print name of person mailing paper or fee)</p> <p><u>Mirella Ponce</u> (Signature of person mailing paper or fee)</p> |
|---|

BOX PCT

Assistant Commissioner for Patents
Washington, D.C. 20231

To the United States Designated Office (DO/US):

- I. Accompanying this transmittal letter are certain items which are required under 35 U.S.C. 371 in order that United States National processing of the above identified International application may commence:
- (X) at the expiration of the applicable time limit under PCT Articles 22 and 39(1) according to the provisions of 35 U.S.C. 371(b).
- () as soon as possible upon receipt of this express request under 35 U.S.C. 371(f).

510 Rec'd FICT/PTO 28 JUL 1999

1. The U.S. National fee [35 U.S.C. 371(c)(1)]

a. () was previously transmitted by applicant on (date)_____.

b. (X) is submitted herewith as follows:

| <u>FOR</u> | <u>NO. FILED</u> | <u>NO. EXTRA</u> | <u>SMALL ENTITY</u> | | | <u>OTHER THAN</u> | |
|---|----------------------------|------------------|---------------------|------------|-----------|-------------------|------------|
| | | | <u>RATE</u> | <u>FEE</u> | <u>or</u> | <u>RATE</u> | <u>FEE</u> |
| Basic Fee | (USPTO NOT ISA OR IPEA) | | //// | \$485 | <u>or</u> | ///// | \$970 |
| Total Claims | - 20 = | -- | x 9 = | | <u>or</u> | x18 = | |
| Ind. Claims | 1 - 3 | -- | x39 = | | <u>or</u> | x78 = | |
| (X) Multiple Dependent Claim Presented | | | +130 = | | <u>or</u> | +260 = | 260 |
| | <u>TOTAL</u> | | | | | | |
| | <u>NATIONAL FEE</u> | | | \$_____ | <u>or</u> | | \$1270 |

i. () A check in the amount of \$_____ is enclosed.

ii. (X) Please charge the filing fee, multiple dependent claim fee (if applicable), excess independent claims fee (if applicable), and excess total claims fee (if applicable) to **Deposit Account No. 23-1703**.

iii. (X) The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 23-1703. A duplicate copy of this sheet is enclosed.

(iv) () The filing fee is not enclosed.

2. A copy of the International application as filed [35 U.S.C. 371(c)(2)]:

a. (X) is transmitted herewith.

b. () is not required as the application was filed with the United States Receiving Office.

c. () has been transmitted

- i. ☐ by the International Bureau. Date of mailing of the application (from form PCT/IB/308): _____ A copy of form PCT/IB/308 is enclosed.
- ii. ☐ by applicant on (date) _____.
3. A translation of the International application into the English language [35 U.S.C. 371(c)(2)]:
- a. ☐ is transmitted herewith.
- b. ☒ is not required as the application was filed in English.
- c. ☐ was previously transmitted by applicant on (date) _____.
4. Amendments to the claims of the International application under PCT Article 19 [35 U.S.C. 371(c)(3)]:
- a. ☐ are transmitted herewith.
- b. ☐ have been transmitted
- i. ☐ by the International Bureau. Date of mailing of the amendments (from form PCT/IB/308): _____.
- ii. ☐ by applicant on (date) _____.
- c. ☒ have not been transmitted as
- i. ☐ no notification has been received that the International Searching Authority has received the Search Copy.
- ii. ☐ the Search Copy was received by the International Searching Authority but the Search Report has not yet issued. Date of receipt of Search Copy (from form PCT/ISA/202): _____.
- iii. ☐ applicant chose not to make amendments under PCT Article 19. Date of mailing of Search Report (from form PCT/ISA/210): _____.

510 Rec'd PCT/PTO 28 JUL 1999

- iv. (X) the time limit for the submission of amendments has not yet expired. The amendments or a statement that amendments have not been made will be transmitted before the expiration of the time limit under PCT Rule 46.1.
5. A Translation of the amendments to the claims under PCT Article 19 [35 U.S.C. 371(c)(3)]:
- a. () is transmitted herewith.
 - b. () is not required as the amendments were made in the English language.
 - c. (X) has not been transmitted for reasons indicated at point I.4.b. or c. above.
6. A declaration for patent application of the inventor [35 U.S.C. 371(c)(4)] complying with 35 U.S.C. 115:
- a. () was previously submitted by applicant on (date)

 - b. (X) is submitted herewith;
and such oath or declaration
 - i. (X) is attached to the application.
 - ii. (X) identifies the application and any amendments under PCT Article 19 which were transmitted as stated in points 1.2.b. or c. and 1.4. and states that they were reviewed by the inventor as required by 37 CFR 1.70.
 - c. () will be submitted subsequently.

II. Concerning other documents:

1. An International Search Report or Declaration under PCT Article 17(2)(a):
- a. () has been transmitted by the International Bureau. Date of mailing (from form PCT/IB/308): _____ A copy of form PCT/IB/308 is enclosed
 - b. () is not required as the application was searched by the United States International Searching Authority.

c. ☐ A copy of the International Search Report is transmitted herewith.

d. ☐ has been submitted by applicant on (date) _____.

2. A Statement of prior art under 37 CFR 1.97 and 1.98:

a. ☐ is transmitted herewith including copies of the references cited on the attached form PTO-1449. Also included is a copy of the International-Type Search Report issued in the Swedish priority document.

b. ☐ will be transmitted within THREE MONTHS of the date of submission of requirements under 35 U.S.C. 371(c).

c. ☐ was previously submitted by applicant on _____, in application serial no. _____.

3. ☒ An Assignment is transmitted herewith for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.

a. ☒ Please charge the \$40 assignment recordation fee to Deposit Account No. 23-1703.


b. ☐ A check in the amount of \$___ is enclosed.

4. Other document(s) or information included:

- Copy of PCT/RO/101 - The PCT Request Form;
- 8 sheets of drawings;
- hard copy of the sequence listing (computer readable form not submitted at this time); and
- a return postcard.

July 28, 1999
DATE

Respectfully submitted,


Richard J. Sterner
Reg. No. 35,372

White & Case LLP
Patent Department
1155 Avenue of the Americas
New York, NY 10036-2787

8/PRTS

510 Rec'd FICT/PTO 28 JUL 1999

1

RNA POLYMERASE ASSAY

TECHNICAL FIELD

5 The present invention relates to a method of identifying a ligand, in particular an inhibitor, of a bacterial RNA polymerase sigma subunit.

BACKGROUND ART

10

Sigma subunits of RNA polymerase

Transcription of genes to the corresponding RNA molecules is a complex process which is catalyzed by DNA dependent RNA polymerase, and involves many
15 different protein factors. In eubacteria, the core RNA polymerase is composed of α , β , and β' subunits in the ratio 2:1:1. To direct RNA polymerase to promoters of specific genes to be transcribed, bacteria produce a variety of proteins, known as sigma (σ) factors, which interact with RNA polymerase to form an active holoenzyme. The resulting complexes are able to recognize and attach to selected
20 nucleotide sequences in promoters.

Physical measurements have shown that the sigma subunit induces conformational transition upon binding to core RNA polymerase. Binding of the sigma subunit to the core enzyme increases the binding constant of the core
25 enzyme for DNA by several orders of magnitude (Chamberlin, M.J. (1974) Ann. Rev. Biochem. 43, 721-).

Bacterial sigma factors do not have any homology with eukaryotic transcription factors, and are consequently a potential target for antibacterial compounds.

Mutations in the sigma subunit, effecting its association and ability to confer DNA sequence specificity to the enzyme, are known to be lethal to the cell.

Characterization of sigma subunits, identified and sequenced from various organisms, allows them to be classified into three groups. The Group I sigma has also been referred to the sigma⁷⁰ class, or the "house keeping" sigma group (for a review see Lonetto et al. (1992) J. Bacteriol. 174, 3843-3849). Sigma subunits belonging to this group recognize similar promoter sequences in the cell. These properties are reflected in certain regions of the proteins which are highly conserved between species.

Another important feature of sigma-dependent transcription is that the sigma subunit dissociates from the core enzyme during elongation of mRNA. Consequently, molecules which stabilize the interaction of the sigma subunit with the core enzyme would also be capable of inactivating transcription.

Mycobacterium tuberculosis

Mycobacterium tuberculosis is a major pulmonary pathogen which is characterized by its very slow growth rate. As a pathogen it gains access to the alveolar macrophages where it multiplies within the phagosome, finally lysing the cells and being disseminated through the blood stream, not only to other areas of the lung, but also to extrapulmonary tissues.

The pathogen thus multiplies in at least two entirely different environments, involving the utilization of different nutrients and a variety of possible host factors. A successful infection would thus involve the coordinated expression of new sets of genes, transcribed by RNA polymerases associating with different sigma factors. This opens the possibility of targeting not only the sigma⁷⁰ subunits

of *M. tuberculosis*, but also other sigma subunits specific for the different stages of infection and dissemination.

The cloning and expression of *Mycobacterium tuberculosis* *sigA* and *sigB* genes are disclosed in the International Patent Application WO 96/38478 (Astra AB).

Anti-sigma factors

Anti-sigma (Asi) proteins are known in the art. Lysates of bacteriophage T₂ (Khesin et al. (1972) Mol. Gen. Genet 119, 299) or phage T₄ (Bogdanova et al. (1970) Mol. Biol. 4, 435; Stevens, A. (1972) Proc. Natl. Acad. Sci. U.S.A. 69, 603) have been reported to inhibit transcription of bacterial genes.

It has been established that the T₄-dependent anti-sigma⁷⁰ activity is borne by a 10 kDa protein (Stevens, A. In: RNA Polymerase p. 617-627 (Eds. R. Losick & M. Chamberlin) Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 1976). A 10 kDa protein was shown to co-purify with RNA polymerase from T₄ infected *E. coli* cells and is detached together with sigma⁷⁰ from the core enzyme on phosphocellulose columns.

A gene called *asiA*, coding for the 10 kDa anti-sigma⁷⁰ factor of bacteriophage T₄, has been identified by Orsini et al. (1993) J. Bacteriol. 175, 85-93. The open reading frame encoded a 90 amino acid protein with the deduced sequence
MNKNIDTVRE IITVASILIK FSREDIVENR ANFIAFLNEI GVTHEGRKLN
QNSFRKIVSE LTQEDKKTLL DEFNEGFEGV YRYLEMYTNK (SEQ ID NO: 5). The *asiA*-encoded protein was overproduced in a phage T7 expression system and partially purified. It showed a strong inhibitory activity towards sigma⁷⁰-directed transcription by *E. coli* RNA polymerase holoenzyme. The nucleotide sequence of gene *asiA* has been deposited in the GenBank data base under accession no.

M99441.

Examples of proteins regulating the sigma subunit of RNA polymerase are known also from other systems. An example is the *S. typhimurium* flagellar regulation system, which is a complex system controlled by a set of over 50 genes grouped into 13 flagellar operons. Late operon expression is positively regulated by the *fliA* gene coding for the sigma factor FliA (Suzuki et al. (1978) J. Bacteriol. 133, 904; Suzuki et al. (1981) J. Bacteriol. 145, 1036). On the other hand, the late operons are negatively regulated by the *flgM* gene. A 7.8 kDa protein has been identified as the *flgM* gene product and purified (Ohnishi et al. (1992) Mol. Microbiol. 6, 3149-3157). This FlgM protein was identified as an anti-sigma factor since it was capable to bind the FliA protein and disturbed its ability to form a complex with the RNA polymerase core enzyme.

Similarly, in *B. subtilis* gene expression, the sigma^F factor has been shown to be regulated by a 14 kDa anti-sigma factor encoded by the *spoIIAB* gene (Duncan & Losick (1993) Proc. Natl. Acad. Sci. U.S.A. 90, 2325-2329), while the sigma^B factor is regulated by a 16 kDa anti-sigma factor encoded by the *rsbW* gene (Benson & Haldenwang (1993) Proc. Natl. Acad. Sci. U.S.A. 90, 2330-2334).

The nucleotide sequences of the genes *flgM*, *spoIIAB* and *rsbW* are available in the GenBank data base (Accession Nos.: FLGMST.PRO for *flgM*, SPOIIAB.PRO for *spoIIAB*, M34995.PRO for *rsbW*). The sequences do not show any gross similarity with the *asiA* sequence disclosed by Orsini et al.

WO 96/25170 discloses methods for treating diseases caused by bacterial pathogens by administering the T₄ AsiA protein, which protein is shown to inhibit the RNA polymerase activity of *Bacillus subtilis* and *Mycobacterium smegmatis*.

DISCLOSURE OF THE INVENTION

In accordance with the present invention, there is provided a method of identifying a ligand of a bacterial sigma⁷⁰ subunit which comprises contacting the sigma⁷⁰ subunit or a portion thereof comprising the anti-sigma binding region, with a test compound and a fusion protein of an anti-sigma⁷⁰ factor of bacteriophage T₄, and determining whether the test compound binds competitively with the anti-sigma⁷⁰ factor to the sigma⁷⁰ subunit or portion thereof.

Preferably, the method comprises:

- (i) immobilizing the sigma⁷⁰ subunit or portion thereof on a matrix or solid support;
- (ii) adding the test compound and the fusion protein;
- (iii) adding a first antibody against the fusion protein;
- (iv) adding a labeled second antibody against the first antibody; and
- (v) determining the amount of second antibody bound to the (first antibody - fusion protein - sigma⁷⁰ subunit or portion thereof) complex formed on the matrix or solid support.

The invention provides an *in vitro* method or assay for screening peptide libraries or chemical libraries for compounds (ligands) which mimic the anti-sigma⁷⁰ factor and which thereby have the capability of inhibiting the interaction of core RNA polymerase and sigma⁷⁰ subunits. Such inhibitor compounds are potentially useful in the treatment of bacterial infections.

The capability of the identified ligands to inhibit the interaction between a bacterial core RNA polymerase and a sigma⁷⁰ subunit can be tested by known methods, in particular by a RNA polymerase assay, such as that disclosed in Orsini et al. (*supra*), wherein transcription of a DNA template, such as calf thymus DNA,

T₄ DNA or poly(dA-dT), is measured by the incorporation of a labeled RNA precursor such as [5-³H]UTP.

In the method or assay according to the invention, the sigma⁷⁰ subunit or portion thereof is preferably obtained from *Escherichia coli* (particularly the C-terminal 99 amino acids of the sigma⁷⁰ subunit containing the anti-sigma binding region) or *Salmonella typhimurium*.

In the fusion protein, the anti-sigma⁷⁰ factor of bacteriophage T₄ will preferably have an amino acid sequence as shown in SEQ ID NO: 1 or SEQ ID NO: 2. The fusion protein preferably also comprises glutathione-S-transferase (GST-Asi) or any other suitable marker gene protein product.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1: Map of plasmid vector pARC 8112.

Fig. 2: Map of plasmid vector pARC 8100.

Fig. 3: Map of plasmid vector pARC 8115.

Fig. 4: Map of plasmid vector pARC 8101.

Fig. 5: Map of plasmid vector pARC 8114.

Fig. 6: Map of plasmid vector pARC 8105.

Fig. 7: Map of plasmid vector pARC 8180.

Fig. 8: Activity of GST-Asi protein on sigma⁷⁰-dependent transcription.

Assay: 0.5 µg *E. coli* core polymerase, 2.5 µg sigma⁷⁰ protein, 1.0 µg T₄ DNA and NTPs. Varying concentrations of GST-Asi was preincubated with sigma protein before addition to the reaction mixture. ³H-UTP specific activity was 2800 - 3000 cpm/nmole.

EXAMPLES

- Throughout this description the terms "standard protocols" and "standard procedures", when used in the context of molecular cloning techniques, are to be understood as protocols and procedures found in an ordinary laboratory manual such as: Current Protocols in Molecular Biology, editors F. Ausubel et al., John Wiley and Sons, Inc. 1994, or Sambrook, J., Fritsch, E.F. and Maniatis, T., Molecular Cloning: A laboratory manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY 1989.

EXAMPLE 1: Cloning, expression and purification of *E. coli* sigma⁷⁰

- PCR primers corresponding to the 5'- and 3'-ends of the coding sequences of *E. coli* sigma were designed with the 5'-sequence including a site for the restriction enzyme *Eco*RI and the 3'-sequence including a site for the restriction enzyme *Sal*I. The PCR amplified DNA fragment was restricted with the above mentioned restriction enzymes and cloned into the *Eco*RI-*Sal*I sites of the expression vector pTrc99a (Amann et al. (1988) Gene 69, 301), whereafter the ligation mix was transformed into *E. coli* DH5α. Transformants harboring the recombinant plasmid with the expected restriction profile were identified.

E. coli DH5 α cells harboring the recombinant plasmid, labeled pARC 8112 (Fig. 1), were grown at +37°C in LB till an OD (600 nm) of 0.4 and induced with 1 mM IPTG. More than 50% of the 90 kDa sigma⁷⁰ protein was found as inclusion bodies as had been reported earlier (Borukhov & Goldfarb (1993) Protein expression and purification 4, 503). The overexpressed sigma⁷⁰ was purified and renatured following standard protocols.

The activity of the purified sigma⁷⁰ protein was confirmed by its ability to support transcription mediated by *E. coli* RNA polymerase core enzyme. *E. coli* RNA polymerase core and holoenzyme forms were purified following the protocol of Burgess and Jendriask (1975) Biochemistry 14, 4634-4638. Polymerase activity using T₄ DNA as template was assayed as described by Orsini et al. (1993) J. Bacteriol. 175, 85-93.

EXAMPLE 2: Cloning, expression and purification of *Salmonella typhimurium* sigma⁷⁰

The coding sequence of the sigma⁷⁰ of *S. typhimurium* was amplified using the forward primer shown as SEQ ID NO: 3 and the reverse primer shown as SEQ ID NO: 4. The forward primer includes the site for the restriction enzyme *EcoRI*, while the reverse primer includes the sequence encoding the stop codon and a site for the restriction enzyme *Sall*. *S. typhimurium* DNA was used as template and the amplified fragment digested with appropriate enzymes. This fragment was then ligated to *EcoRI* - *Sall* digested pTrc 99A (Amann et al. (1988) Gene 69, 301) and transformed into *E. coli* DH5 α to obtain the recombinant plasmid pARC 8118. Plasmid DNA of pARC 8118 was digested with *EcoRI* - *HindIII* and the released 2.3 kb DNA fragment encompassing the entire coding sequence of *S. typhimurium* sigma⁷⁰ was cloned into the *EcoRI* - *HindIII* digested pRSET B (Kroll et al. (1993) DNA and Cell Biol. 12, 441) vector which has not only

the T7 promoter but also sequences encoding a his tag which is now fused at the C-terminus of *S. typhimurium* sigma⁷⁰. The recombinant plasmid encoding the *S. typhimurium* sigma⁷⁰ fused to a His tag was labeled pARC 8133.

5 Plasmid DNA of pARC 8133 was then used to transform the expression host *E. coli* BL21(DE3) and transformants obtained at +37°C. Cells harboring pARC 8133 were grown in LB at +37°C till an OD of 0.4 at 600 nm and induced with 1 mM IPTG for 4 hours. The cells were pelleted, suspended in buffer and sonicated, according to standard methods. The lysed cells were clarified by centrifugation at 45,000 rpm
10 for 60 min at +4°C and the clarified supernatant was loaded directly on Ni-agarose (Pharmacia) and washed with 50 mM imidazole. The bound protein was eluted with 200 mM imidazole and dialysed overnight at +4°C against 50 mM Tris-Cl, pH 7.5. The eluate was concentrated using Amicon cone filters to 1/10 volume. The concentrated protein was cleaved with enterokinase, used at a ratio of 1:50 of
15 the enzyme, at +25°C for 14 hours. The cleaved mixture was passed through a Ni²⁺ agarose column and the unbound material collected. The cleaved protein was analyzed for homogeneity by SDS-PAGE.

20 EXAMPLE 3: Toxicity of the anti-sigma protein

3.1. Cloning of the *asiA* gene

The coding sequence for the *asiA* gene (Orsini et al., *supra*) was amplified from the
25 genomic DNA of bacteriophage T₄ by PCR using the forward primer shown as SEQ ID NO: 5 and the reverse primer designated shown as SEQ ID NO: 6.

The 5'-primer included the sequence of the restriction enzyme *NcoI* while the 3'-primer included the sequence for the restriction enzyme *BamHI*. The coding
30 sequence was amplified by PCR following standard protocols and the amplified

fragment ligated to *Nco*I - *Bam*HI restricted pBR329 and transformed into competent cells *E. coli* DH5 α . Recombinants were selected at +37°C as chloramphenicol sensitive, ampicillin resistant transformants. One of the transformants having the desired restriction pattern was labeled pARC 8100 (Fig. 2).

The inclusion of the sequence encoding *Nco*I resulted in the change of the second amino acid from asparagine to glycine (SEQ ID NO: 2). The nucleotide sequence of the *asiA* gene cloned in pARC 8100 was verified by double stranded sequencing and found to be identical to the *asiA* sequence as disclosed by Orsini et al., except for the expected change of codon for the second amino acid as a result of the PCR cloning protocol used.

3.2. *In vivo* toxicity of *asiA* product in *E. coli*

(A)

The *Nco*I - *Bam*HI DNA fragment from pARC 8100 was ligated to pET 8c Km (Umender K. Sharma et al., J. Bact., 177, 6745 (1996)) and kanamycin resistant transformants with *E. coli* DH5 α were selected at +37°C. One of the transformants harboring a plasmid with the expected restriction enzyme profile was labeled pARC 8115 (Fig. 3). pARC 8115 plasmid DNA was then used to transform the expression host *E. coli* BL21(DE3) (Studier et al., *supra*) and transformants selected both at +37°C and at +30°C. However, no viable transformants could be obtained at either of the temperatures. The leaky expression from the T7 promoter in pET 8c being much higher than from pET 11d(Km) (Studier et al., *supra*), the toxicity of the *asiA* product could explain the non-transformability.

Transformants could however be obtained using *E. coli* BL21(DE3)/pLysS as host, in which the leaky expression is additionally repressed by the T7 lysozyme expressed from pLysS.

(B)

The *Nco*I - *Bam*HI 284 bp DNA fragment was obtained from pARC 8100 and ligated to the *Nco*I - *Bam*HI sites of pET 11d Km (Umender K. Sharma et al., J. Bact., 177, 6745 (1996)) and transformed into *E. coli* DH5 α . Transformants were selected for kanamycin resistance. Plasmid preparations from individual transformants were digested with restriction enzymes and the correct transformant that released the 284 bp fragment after *Nco*I - *Bam*HI digest was labeled pARC 8101 (Fig. 4). The transformants were selected at +37°C and appeared normal.

pARC 8101 DNA was then used to transform the expression host *E. coli* BL26(DE3). BL26 is a lac Iq isogenic strain of BL21 (Studier et al., *supra*). Transformants selected for kanamycin both at +37°C and at +30°C.

The transformants obtained at +37°C were morphologically sick and non-viable indicating the acute toxicity of the leaky expression of the *asiA* gene. In contrast, healthy colonies could be obtained when transformants were selected at +30°C where the leaky expression from the T7 promoter can be expected to be negligible.

BL26(DE3)/pLysS colonies were healthy both at +37°C and +30°C, indicating that the tight regulation of expression in this strain made the *asiA* gene non-toxic to the host. Transformants obtained from DH5 α were also healthy both at +37°C and +30°C, indicating that the gene when transformed into a non-expression host was non-toxic.

(C)

The *asiA* gene was excised from pARC 8101 as a *Xba*I - *Bam*HI DNA fragment to include the sequence for the ribosome binding site and ligated to the low copy vector, *Xba*I - *Bam*HI cleaved pWKS129 (Wang & Kushner (1991) Gene 100, 195) and the ligation mix transformed to *E. coli* DH5 α . Recombinant plasmid harboring the *asiA* gene was identified by restriction profile and labeled pARC 8114 (Fig. 5).

This plasmid DNA when transformed into *E. coli* BL21(DE3) gave viable transformants both at +37°C and +30°C, since the low copy number of the plasmid reduced the level of the AsiA protein expressed through leaky expression.

5

EXAMPLE 4: Glutathione-S-transferase - Anti-sigma fusion protein

4.1. Cloning of fusion protein in *E. coli*

- 10 The coding sequence for the *asiA* gene was excised as an *Nco*I - *Bam*HI fragment from pARC 8100 and ligated to *Nco*I - *Bam*HI cleaved pARC 0499. The plasmid pARC 0499 has a *Nco*I site in frame with the glutathione-S-transferase encoding sequence, enabling fusion of the N-terminus of *asiA* sequences. The ligation mix was transformed into *E. coli* DH5 α and transformants selected at +37°C and
- 15 +30°C. Viable colonies were obtained at both temperatures indicating the N-terminal fusion reduces the toxicity of the protein. The recombinant plasmid obtained with the sequences encoding GST-AsiA was labeled pARC 8105 (Fig. 6).

- The GST-AsiA fusion protein was purified as follows: *E. coli* DH5 α harboring
- 20 pARC 8105 was grown in LB till an OD at 600 nm of 0.6 at +37°C followed by the addition of 1 mM IPTG and allowing growth for a further period of 2 hours. The cells were then centrifuged at 5000 rpm for 10 minutes. The pelleted cells were suspended in Buffer A (phosphate buffered saline (PBS) pH 7.5, 5 μ g/ml aprotinin, 5 μ g/ml leupeptin) and sonicated. The sonicate was clarified at 45,000 rpm for 10
- 25 minutes at +4°C and the supernatant passed through a Glutathione-Sepharose 4B column previously equilibrated with PBS. The column was washed with 5 bed volumes of 1 M urea once followed by washing with PBS (10 bed volumes). The bound protein was eluted with 10 mM glutathione and the eluate dialysed overnight against buffer containing 50 mM Tris(hydroxymethyl)aminomethane

hydrochloride (Tris HCl), pH 7.5. The dialysed protein was then concentrated to 1/10 vol using Amicon cone filters. The concentrated protein was then treated with 1:50 ratio of Factor Xa protease in a buffer containing 50 mM Tris HCl, pH 7.5, 100 mM NaCl, 1 mM CaCl_2 at +25°C for 4 hours. The digested eluate was then
5 passed through a Glutathione-Sepharose column and the unbound fraction collected and analyzed for the purity of the AsiA protein by SDS-PAGE.

4.2. Cloning of fusion protein in yeast

10 The expression and purification of the GST-AsiA fusion protein in *E. coli* did not allow large scale production because of the inherent toxicity of the fusion protein. To obtain large quantities of the AsiA protein, the possibility of expressing the GST-AsiA fusion protein in a yeast host was investigated. As the *Saccharomyces* species do not have a sigma⁷⁰ homologue, the AsiA protein was expected not to
15 have toxicity effects on the transcription apparatus of the host and thus for initial experiments a *Saccharomyces cerevisiae* expression system was chosen.

The gene encoding the GST-AsiA fusion protein on the plasmid pARC 8105 was amplified using the forward primer shown as SEQ ID NO: 7 and the reverse
20 primer shown as SEQ ID NO: 8. The amplified 1.0 kb fragment was restricted with *Bgl*III and *Hind*III which are the sites introduced by the primer sequence at the 5'- and 3'-ends of the amplified PCR fragment. The *Bgl*III – *Hind*III restricted fragment was ligated into the *Bgl*III – *Hind*III restricted *Saccharomyces cerevisiae* cloning vector pSW6 (Pascall et al. (1991) J. Mol. Endocrinol. 6, 63-70). The ligation mixture
25 was transformed into the *Saccharomyces cerevisiae* host CGY1585 and transformants selected on leucine (3 µg/ml).

Individual transformants from both reaction mixtures were screened for the presence of plasmid with the expected restriction profile. The recombinant
30 plasmid obtained with the vector PSW6 was labeled pARC 8180 (Fig. 7).

S. cerevisiae CGY 1585 strain harbouring pARC 8180 were grown using standard procedures and lysed by passing the concentrated cell suspension through a French Press. The lysate was then centrifuged at 10,000 rpm and the supernatant applied to a Glutathione-Sepharose column (Pharmacia) and the fusion protein
5 purified following standard protocols.

The purified GST-AsiA fusion protein was cleaved with Factor Xa as recommended and the AsiA protein separated. The yield of the purified AsiA protein was 500 µg/l. The activity of the GST-AsiA protein, expressed and purified
10 from *Saccharomyces*, was compared to that obtained from *E. coli* and found to be identical in its ability to inhibit sigma⁷⁰ mediated transcription of T₄ template (Fig. 8). AsiA protein purified from GST-AsiA after Factor Xa cleavage also inhibited *E. coli* sigma⁷⁰ dependent transcription.

15 The ability to overexpress and purify large quantities of the AsiA protein in *Saccharomyces* without toxicity problems for the host also indicates that the AsiA protein does not have a corresponding sigma⁷⁰ homologue in this species. As the *Saccharomyces* RNA polymerase is similar to that of higher eukaryotes it also substantiates the fact that the AsiA protein is toxic only to the prokaryotic
20 transcription apparatus.

In order to test the feasibility of expressing AsiA in alternative yeast expression systems, *Pichia pastoris* expression system was chosen. The gene encoding GST-AsiA was amplified by PCR from pARC 8105 and cloned into pPIC9K
25 (Invitrogen).

5' primer (asi34) - TA TAC GTA TCC CCT ATA CTA GGT TAT TGG

3' primer (asi35) - T TGC GGC CGC TTA TTT GTT CGT ATA CAT

PCR conditions used were:

| | |
|-----------------------|-------|
| melting temperature | 94 °C |
| annealing temperature | 50 °C |
| extension temperature | 72 °C |

and 30 cycles were performed.

1Kb PCR product was cloned as SnaBI-NotI fragment into pPIC9K (pARC 8274).

Pichia pastoris transformants of pARC 8274 were selected based on G 418 resistance. Expression studies indicated that GST-AsiA was secreted into the culture supernatant. GST-AsiA purified from the culture supernatant was found to inhibit *in vitro* transcription as efficiently as that of either *E. coli* or the *Saccharomyces* GST-AsiA.

This indicates that it is feasible to produce recombinant GST-AsiA in a yeast expression system.

EXAMPLE 5: *In vitro* AsiA-sigma⁷⁰ interaction assays

5.1. RNA polymerase assay

The *E. coli* RNA polymerase assay was standardized following the protocol of Orsini et al. (J. Bacteriol. 175, 85-93, 1993) using T₄ phage DNA as template to quantify sigma⁷⁰ dependent transcription. *E. coli* RNA polymerase core enzyme was purified following the protocol of Burgess and Jendriask (1975) Biochemistry 14, 4634-4638.

5.2. Inhibition of *E. coli* transcription by AsiA protein

Increasing concentrations of AsiA protein expressed in *S. cerevisiae* was added to the reaction mixture and the transcription mediated by the *E. coli* RNA polymerase
5 quantified. The results showed that increasing concentrations of AsiA protein completely inhibited transcription mediated by *E. coli* sigma⁷⁰.

5.3. Inhibition of *E. coli* transcription by GST-AsiA fusion protein

10 The GST-AsiA fusion protein was purified from *S. cerevisiae* as described in Section 5.2 and used at different concentrations in the *E. coli* sigma⁷⁰ mediated RNA polymerase transcription assay, using phage T₄ DNA as template. The GST-AsiA fusion protein inhibited >80% the sigma⁷⁰ mediated transcription when the core enzyme was reconstituted with 2.5 µg of sigma⁷⁰ protein.

15

5.4. Reactivation of *E. coli* sigma⁷⁰ mediated transcription

Increasing concentrations of *E. coli* sigma⁷⁰ protein were added to the reaction mixture containing 0.1 µg of AsiA. As shown in Table 1, addition of sigma⁷⁰
20 protein could reactivate transcription mediated by *E. coli* RNA polymerase. This reversal of inhibition demonstrates the specific interaction of the AsiA protein with sigma⁷⁰.

TABLE 1

| | <i>E. coli</i> sigma ⁷⁰ | <i>S.</i> <i>typhimurium</i> sigma ⁷⁰ |
|---|---------------------------------------|--|
| | [³ H]UTP (nmol) | |
| <i>E. coli</i> RNA polymerase | 2.1 | 2.3 |
| <i>E. coli</i> RNA polymerase + 0.1 µg AsiA | 0.57 | 0.34 |
| <i>E. coli</i> RNA polymerase + 0.1 µg AsiA + 4 µg sigma ⁷⁰ | 2.3 | 3.0 |
| <i>E. coli</i> RNA polymerase + 0.1 µg AsiA + 4 µg sigma ⁷⁰ (AsiA preincubated) | 0.68 | 0.98 |

5.5. Effect of *Salmonella typhimurium* sigma⁷⁰ on *E. coli* transcription

As described in Section 5.1, the *E. coli* RNA polymerase activity assay was standardized with purified enzyme and phage T₄ DNA as template. The RNA polymerase could be >80% inhibited by 0.07 µg of purified AsiA protein.

- 10 To a reaction mixture containing the *E. coli* RNA polymerase, T₄ DNA and 0.1 µg of AsiA, increasing concentrations of purified *Salmonella typhimurium* sigma⁷⁰ was added and the RNA polymerase activity quantified. As shown in Table 1, addition of 4 µg of *S. typhimurium* sigma⁷⁰ could restore the activity of the RNA polymerase.

15

5.6. Preincubation of sigma⁷⁰ with AsiA

To a reaction mixture containing *E. coli* RNA polymerase and T₄ DNA template there was added 0.1 µg AsiA and 4 µg *E. coli* or *S. typhimurium* sigma⁷⁰,

preincubated at +37°C for 10 min with 0.1 µg of AsiA. As shown in Table 1, preincubation of the sigma⁷⁰ with AsiA abolished the ability of the sigma⁷⁰, from both *E. coli* and *S. typhimurium*, to activate *E. coli* RNA polymerase.

5

EXAMPLE 6: Competitive ELISA for quantification of AsiA-sigma⁷⁰ interaction

Screening assays

10 The above principles provide a basis for a method of identification of oligonucleotide sequences encoding peptides, which may be either related or unrelated to the AsiA peptide, and which could efficiently bind to sigma⁷⁰ subunits from *E. coli*, *S. typhimurium* and/or other housekeeping or virulence-associated sigma subunits of bacterial pathogens.

15

The identification of such peptide structures would enable the further identification, by known methods, of peptoids, peptidomimetics and organic molecules which can be tested in transcription assays for inhibition of sigma⁷⁰ dependent transcription.

20

Truncated sigma⁷⁰ (C-terminal 99 amino acids) which has been shown to be the anti-sigma binding region of sigma⁷⁰ of *E. coli* (Severinova et al., J. Mol. Biol. (1996) 263 (5), pp 637-647) was cloned and expressed in a His tag expression vector resulting in a (His)₅-tagged protein product.

25

A solution of the protein product (200 pmoles/ml) in a pH 7.5 buffer comprising Tris HCl (10 mM) and sodium chloride (50 mM) was prepared and 200 µl of solution were added per well of a nickel-coated microtitre plate, followed by incubation at room temperature for a period of two hours. The microtitre plate
30 was then washed 5 times with the buffer solution and then incubated at 37°C for

one hour with a solution of bovine serum albumin (3% w/v) in the above buffer. After washing 5 times with phosphate buffered saline (PBS) with "Tween" 20 (polyoxyethylene (20) sorbitan monolaurate, 0.5% v/v), the GST-AsiA fusion protein prepared as described in *Example 4.1.* above (100 ng) was added together
 5 with a defined amount of a putative inhibitor of truncated sigma⁷⁰ protein (corallopyronin, myxopyronin or ripostatin) and incubated at 37°C for one hour. Further washing (5 times) with PBS-"Tween" 20 was carried out and then anti-GST antibodies (1:2000) raised in rabbits were added to the wells, with incubation at 37°C for one hour. The previous washing step was repeated and thereafter anti-
 10 rabbit IgG-horse radish peroxidase (HRP) conjugate (1:2000) was added and incubated at 37°C for one hour. A final wash with PBS-"Tween" 20 was carried out, before adding the HRP enzyme substrate, tetramethyl benzidine/ H₂O₂. The enzymic reaction (resulting in the development of colour) was stopped after a suitable period of time by the addition of 6N H₂SO₄ and the microtitre plate was
 15 "read" in a spectrophotometer using light of wavelength (λ) 450 nm. The results obtained are shown in Table 2 following.

TABLE 2

| Test | % Binding to truncated sigma ⁷⁰ |
|---|--|
| GST-AsiA (500 ng/ml) | 100 |
| GST-AsiA + AsiA (7 μ g/ml) | 12 |
| GST-AsiA + corallopyronin (30 μ g/ml) | 69 |
| GST-AsiA + myxopyronin (30 μ g/ml) | 56 |
| GST-AsiA + ripostatin (1 μ g/ml) | 100 |
| GST-AsiA + ripostatin (10 μ g/ml) | 55 |
| GST-AsiA + ripostatin (100 μ g/ml) | 1.5 |

CLAIMS

1. A method of identifying a ligand of a bacterial sigma⁷⁰ subunit which comprises contacting the sigma⁷⁰ subunit or a portion thereof comprising the anti-sigma binding region, with a test compound and a fusion protein of an anti-sigma⁷⁰ factor of bacteriophage T₄, and determining whether the test compound binds competitively with the anti-sigma⁷⁰ factor to the sigma⁷⁰ subunit or portion thereof.
2. A method according to claim 1, which comprises:
 - (i) immobilizing the sigma⁷⁰ subunit or portion thereof on a matrix or solid support;
 - (ii) adding the test compound and the fusion protein;
 - (iii) adding a first antibody against the fusion protein;
 - (iv) adding a labeled second antibody against the first antibody; and
 - (v) determining the amount of second antibody bound to the (first antibody - fusion protein - sigma⁷⁰ subunit or portion thereof) complex formed on the matrix or solid support.
3. A method according to claim 1 or claim 2, wherein the sigma⁷⁰ subunit or portion thereof is obtained from *Escherichia coli* or *Salmonella typhimurium*.
4. A method according to any one of the preceding claims, wherein the anti-sigma⁷⁰ factor has an amino acid sequence as shown in SEQ ID NO: 1 or SEQ ID NO: 2.
5. A method according to any one of the preceding claims, wherein the fusion protein also comprises glutathione-S-transferase.

6. A method according to any one of the preceding claims wherein the ligand is an inhibitor of a bacterial sigma⁷⁰ subunit.

A B S T R A C T

RNA POLYMERASE ASSAY

- 5 The invention provides a method of identifying a ligand, in particular an inhibitor, of a bacterial RNA polymerase sigma subunit.




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. B

DECLARATION FOR PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled RNA POLYMERASE ASSAY the specification of which is attached hereto unless the following box is checked:

☒ was filed on 07 June 1999 as United States Application Number or PCT International Application Number SE99/00979 and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Priority Not Claimed

| | | | |
|-------------------------|---------------------|--|--------------------------|
| 1239/MAS/98 (Number) | India (Country) | 09 June 1998 (Day/Month/Year Filed) | <input type="checkbox"/> |
| 9802573-7 (Number) | Sweden (Country) | 17 July 1998 (Day/Month/Year Filed) | <input type="checkbox"/> |

I hereby claim the benefit under 35 U.S.C. § 119(e) of any United States provisional application(s) listed below.

| | |
|----------------------|---------------|
| (Application Number) | (Filing Date) |
| (Application Number) | (Filing Date) |

I hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. § 112.

I acknowledge the duty to disclose information which is material to patentability as defined in 37 CFR § 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application.

| | | |
|----------------------|---------------|--|
| (Application Number) | (Filing Date) | (Status -- patented, pending, abandoned) |
| (Application Number) | (Filing Date) | (Status -- patented, pending, abandoned) |

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

(13) Edward V. Filardi, Reg. No. 25,757; Nels T. Lippert, Reg. No. 25,888; Dimitrios Drivas, Reg. No. 32,218; Robert B. Smith, Reg. No. 28,538; Cecilia O'Brien Lofters, Reg. No. 33,434; David Bender, Reg. No. 35,445; John M. Genova, Reg. No. 32,224; Richard J. Sterner, Reg. No. 35,372; Hans-Peter G. Hoffmann, Reg. No. 37,352; Thelma C. Cleland, Reg. No. 40,948; Leslie Morioka, Reg. No. 40,304; John Scheibeler, Reg. No. 35,346; and Roy Waldron III, Reg. No. 42,208, all of the firm of WHITE & CASE Limited Liability Partnership, with offices at 1155 Avenue of the Americas, New York, New York 10036.

Address all telephone calls to _____ at telephone number (212) 819-8200

Address all correspondence to WHITE & CASE LLP
Patent Department
1155 Avenue of the Americas
New York, NY 10036-2787

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believe to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor
(given name, family name)

Tanjore Balganes

First inventor's signature

Tanjore Balganes

Date:

15 June 1995

Residence Address

Bangalore, India

INX

Citizenship

Indian

Post Office Address

Astra Biochemicals Pvt Ltd, P.O. Box 8013, Malleswaram, Bangalore 560080, India

Full name of second joint inventor, if any ²⁻⁰⁰
(given name, family name)

Vasanthi Ramachandran

Second inventor's signature

Ramachandran

Date

15 June 1999

Residence Address

Bangalore, India

INX

Citizenship

Indian

Post Office Address

Astra Biochemicals Pvt Ltd, P.O. Box 8013, Malleswaram, Bangalore 560080, India

Full name of third joint inventor, if any ³⁻⁰⁰
(given name, family name)

Umender Sharma

Third inventor's signature

Umender

Date

15 June 1999

Residence Address

Bangalore, India

INX

Citizenship

Indian

Post Office Address

Astra Biochemicals Pvt Ltd, P.O. Box 8013, Malleswaram, Bangalore 560080, India

Full name of fourth joint inventor, if any
(given name, family name)

Fourth inventor's signature

Date

Residence Address

Citizenship

Post Office Address

Full name of fifth joint inventor, if any
(given name, family name)

Fifth inventor's signature

Date

Residence Address

Citizenship

Post Office Address

Additional inventors are being named on separately numbered sheets attached hereto.

SEQUENCE LISTING

<110> ASTRA AKTIEBOLAG

<120> RNA Polymerase Assay

<130> R 1765

<140>

<141>

<160> 8

<170> PatentIn Ver. 2.0

<210> 1

<211> 90

<212> PRT

<213> Bacteriophage T4

<400> 1

Met Asn Lys Asn Ile Asp Thr Val Arg Glu Ile Ile Thr Val Ala Ser
1 5 10 15

Ile Leu Ile Lys Phe Ser Arg Glu Asp Ile Val Glu Asn Arg Ala Asn
20 25 30

Phe Ile Ala Phe Leu Asn Glu Ile Gly Val Thr His Glu Gly Arg Lys
35 40 45

Leu Asn Gln Asn Ser Phe Arg Lys Ile Val Ser Glu Leu Thr Gln Glu
50 55 60

Asp Lys Lys Thr Leu Ile Asp Glu Phe Asn Glu Gly Phe Glu Gly Val
65 70 75 80

Tyr Arg Tyr Leu Glu Met Tyr Thr Asn Lys
85 90

<210> 2

<211> 90

<212> PRT

<213> Bacteriophage T4

<400> 2

Met Gly Lys Asn Ile Asp Thr Val Arg Glu Ile Ile Thr Val Ala Ser
1 5 10 15

Ile Leu Ile Lys Phe Ser Arg Glu Asp Ile Val Glu Asn Arg Ala Asn
 20 25 30

 Phe Ile Ala Phe Leu Asn Glu Ile Gly Val Thr His Glu Gly Arg Lys
 35 40 45

 Leu Asn Gln Asn Ser Phe Arg Lys Ile Val Ser Glu Leu Thr Gln Glu
 50 55 60

 Asp Lys Lys Thr Leu Ile Asp Glu Phe Asn Glu Gly Phe Glu Gly Val
 65 70 75 80

 Tyr Arg Tyr Leu Glu Met Tyr Thr Asn Lys
 85 90

<210> 3
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 3
 atggaattca acccgcagtc acagctg 27

<210> 4
 <211> 27
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 4
 tgagtcgact taatcgtcga ggaagct 27

<210> 5
 <211> 25
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: PCR primer

<400> 5

ggccatgggc aataaaaaaca ttgat

25

<210> 6

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 6

ggggatcctt atttggtcgt atacat

26

<210> 7

<211> 30

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 7

gaaagatctc atatgtcccc tatactaggt

30

<210> 8

<211> 29

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 8

ctaagctttt atttggtcgt atacatctc

29